



ON YOUR SIDE

For Extensive-Stage Small Cell Lung Cancer (ES-SCLC)

Your Guide

WHAT IS TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat adults with a **type of lung cancer called small cell lung cancer (SCLC)**. **TECENTRIQ may be used with the chemotherapy medicines carboplatin and etoposide as your first treatment when your lung cancer is a type called "extensive-stage small cell lung cancer,"** which means that it has spread or grown.

It is not known if TECENTRIQ is safe and effective in children.

SELECT IMPORTANT SAFETY INFORMATION

TECENTRIQ can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life threatening and can lead to death.

Please see pages 20-21 and full [Prescribing Information](#) for additional Important Safety Information.

This guide will help explain how TECENTRIQ is thought to work and will provide Important Safety Information. This brochure also includes a list of financial resources that might be available. Read up on some practical treatment tips that may be helpful for you and your support circle.

This brochure is not intended to replace a conversation with your doctor.

Here for you

Dealing with cancer can be difficult and unexpected. It's okay to feel overwhelmed or unsure of what the future looks like. There may be ups and downs ahead. This guide is here to help you navigate your journey and provide you with support. Learn more about extensive-stage small cell lung cancer (ES-SCLC) and how TECENTRIQ may help. This is a great starting point to help you prepare for the next steps ahead.

For more information visit



[TECENTRIQ.com/SCLCinfo](https://www.tecentriq.com/SCLCinfo)

or call the Patient Resource Center:



1 (877) GENENTECH (1-877-436-3683)

Monday - Friday, 6AM - 5PM PT

The Patient Resource Center is a general help line for questions related to TECENTRIQ for patients and caregivers and is not intended to be medical advice or replace the advice of your healthcare team.

Getting Started

This brochure was made to help answer questions you may have about extensive-stage small cell lung cancer and TECENTRIQ.

3

Remember to talk to your healthcare team throughout the process. They can answer any questions you may have and make sure the treatments you receive are right for you. Find information on the following:

pages **4-5**

What is small cell lung cancer?

What is TECENTRIQ?

pages **6-7**

Understanding cancer cells & PD-L1

How is TECENTRIQ thought to work?

pages **8-9**

What are the potential benefits of TECENTRIQ?

pages **10-11**

**How is TECENTRIQ given?
Financial resources**

pages **12-13**

**Additional financial resources
Connecting with care partners**

pages **15-19**

Information for caregivers

pages **20-21**

**Important Safety Information
about TECENTRIQ**

page **22**

Glossary

Please see full **Prescribing Information** and additional **Important Safety Information** throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE

WHAT IS SMALL CELL LUNG CANCER?

Small cell lung cancer (SCLC) occurs when the cells in your lungs start to grow unusually or uncontrollably. This unusual growth takes place inside your lungs or the lining of the airways. As more cancer cells grow, they can form into a tumor.

SCLC is an aggressive form of cancer because it can grow quickly. When the cancer is first found, it may only be present in one of the two lungs in the chest (limited-stage SCLC). As the disease advances, it may spread to the other lung or to other parts of the body. This is called extensive-stage SCLC. For many people with SCLC, the cancer can spread soon after diagnosis.

WHAT ARE SOME OF THE SYMPTOMS OF SCLC?

The most common symptoms of SCLC may include:

- A cough that does not go away or gets worse
- Coughing up blood or rust-colored spit
- Chest pain that may get worse when taking deep breaths, coughing, or laughing
- Hoarseness
- Weight loss and loss of appetite
- Shortness of breath
- Feeling tired or weak
- Infections such as bronchitis and pneumonia that don't go away or keep coming back
- New onset of wheezing


When small cell lung cancer spreads to distant organs (extensive-stage SCLC), it can cause other symptoms which may include:

- Bone pain (like pain in the back or hips)
- Nervous system changes (such as headache, weakness or numbness of an arm or leg, dizziness, balance problems, or seizures), from cancer spread to the brain
- Yellowing of the skin and eyes (jaundice), from cancer spread to the liver
- Lumps near the surface of the body, due to cancer spreading to the skin or to lymph nodes (collection of immune system cells), such as those in the neck or above the collarbone

Experiencing these symptoms does not necessarily mean you have lung cancer. However, it is very important to talk to your healthcare team if you experience any of these symptoms. Your healthcare team has the knowledge and the tools needed to make the appropriate diagnosis.

ASK YOUR DOCTOR ABOUT TECENTRIQ WITH CHEMOTHERAPY AS A FIRST TREATMENT

TECENTRIQ in combination with chemotherapy (carboplatin and etoposide) can be used as your first treatment for extensive-stage SCLC.



WHAT IS TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat adults with:

A type of lung cancer called small cell lung cancer (SCLC).

• **TECENTRIQ may be used with the chemotherapy medicines carboplatin and etoposide as your first treatment when your lung cancer:**

- is a type called “extensive-stage small cell lung cancer,” which means that it has spread or grown

It is not known if TECENTRIQ is safe and effective in children.

Please see full [Prescribing Information](#) and additional Important Safety Information throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1000 mg
INJECTION FOR IV USE

UNDERSTANDING CANCER CELLS & PD-L1

Your immune system helps to protect you against infection and disease. It does this through a collection of specialized organs and cells. Normally, your immune system is able to detect and remove abnormal cells, such as cancer cells.

6



Some cancer cells can go undetected by your immune system. One way they can go undetected is through a protein called programmed death-ligand 1, or PD-L1. This protein is found on many cells, including cancer cells. PD-L1 on cancer cells may bind to your immune cells. When it does this, the cancer cell is seen as a healthy cell. This may prevent your immune system from being able to recognize and attack your cancer.

HOW IS TECENTRIQ THOUGHT TO WORK?

TECENTRIQ is a cancer immunotherapy that can help reactivate the immune system so it can recognize cancer cells in the body. TECENTRIQ is an anti-PD-L1 antibody, meaning it is designed to block PD-L1. By blocking PD-L1, TECENTRIQ is thought to reveal cancer cells. This process helps your immune system recognize, target, and attack your cancer. TECENTRIQ may also affect normal cells.

SELECT IMPORTANT SAFETY INFORMATION

Possible serious side effects with TECENTRIQ include, but are not limited to, lung problems (pneumonitis), liver problems (hepatitis), intestinal problems (colitis), hormone gland problems (especially the thyroid, adrenal glands, pancreas, and pituitary), problems in other organs, severe infections and severe infusion reactions.



Please see full **Prescribing Information** and additional **Important Safety Information** throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1000 mg
INJECTION FOR IV USE

WHAT ARE THE POTENTIAL BENEFITS OF TECENTRIQ WITH CHEMOTHERAPY?

In a clinical trial of TECENTRIQ with chemotherapy (carboplatin and etoposide) that included **403 people** with ES-SCLC:

MORE PATIENTS LIVED LONGER WITH TECENTRIQ + CHEMOTHERAPY

MEDIAN OVERALL SURVIVAL (OS)*



*Median OS = The length of time from the start of treatment in a trial at which half of the patients are still alive.

- Patients who received TECENTRIQ with chemotherapy had a **30% reduced risk** of dying compared to chemotherapy alone

†Chemotherapy = carboplatin + etoposide.

TECENTRIQ may not work for everyone.

SELECT IMPORTANT SAFETY INFORMATION

TECENTRIQ can cause serious side effects, including:

- **Lung problems (pneumonitis)**-signs and symptoms of pneumonitis may include new or worsening cough, shortness of breath, and chest pain
- **Liver problems (hepatitis)**-signs and symptoms of hepatitis may include yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), drowsiness, dark urine (tea colored), bleeding or bruising more easily than normal, and feeling less hungry than usual

WHAT ARE THE POTENTIAL BENEFITS OF TECENTRIQ WITH CHEMOTHERAPY?

9

In the same clinical trial with **403 people** with ES-SCLC:

PATIENTS HAD DELAYED WORSENING OF THEIR DISEASE WITH TECENTRIQ + CHEMOTHERAPY

MEDIAN PROGRESSION-FREE SURVIVAL (PFS)*



*Median PFS = The length of time from the start of treatment in a trial that half the patients lived without their tumors growing or spreading.

†Chemotherapy = carboplatin + etoposide.

TECENTRIQ may not work for everyone.

Please see full [Prescribing Information](#) and additional
Important Safety Information throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1000 mg
INJECTION FOR IV USE

HOW IS TECENTRIQ GIVEN?

Your healthcare provider will give you TECENTRIQ into your vein through an intravenous (IV) line. They will also decide how long you may be on treatment. If at any point the disease progresses or treatment can't be tolerated, infusions will be stopped.

FIRST 4 INFUSIONS: WITH CHEMOTHERAPY

Every 3 weeks for 12 weeks




TECENTRIQ is given over 60 minutes. If the first infusion is tolerated, all following TECENTRIQ infusions will be given over 30 minutes. Your other medications may take longer.

↓ AFTER 12 WEEKS ↓

LATER INFUSIONS: WITHOUT CHEMOTHERAPY



After completion of chemotherapy, your doctor will continue TECENTRIQ alone. TECENTRIQ alone offers flexible dosing options. Your doctor may suggest one of the following TECENTRIQ infusion schedules:



SPEAK WITH YOUR HEALTHCARE PROVIDER TO CREATE AN INFUSION SCHEDULE FOR YOU

COMMITTED TO HELPING YOU FIND ASSISTANCE OPTIONS FOR TECENTRIQ

If you are worried about paying for TECENTRIQ, we can help you understand options that may help you. We can find out how much it will cost. We can also help you understand if there are options to help you pay for it.

TECENTRIQ ACCESS SOLUTIONS

TECENTRIQ Access Solutions is committed to helping you get the Genentech medicine your doctor prescribed.

At [Genentech-Access.com/TECENTRIQ/patients](https://www.genentech-access.com/TECENTRIQ/patients), you can:

- Find out what we offer
- Learn what to expect
- Get started with patient assistance

GENENTECH PATIENT FOUNDATION

The Genentech Patient Foundation helps people whether or not they have insurance. For more information, call a Foundation Specialist at **(888) 941-3331** or visit [GenentechPatientFoundation.com](https://www.GenentechPatientFoundation.com).

To be eligible for free Genentech medicine from the Genentech Patient Foundation, insured patients who have coverage for their medicine must have exhausted all other forms of patient assistance (including TECENTRIQ Co-pay Program and support from independent co-pay assistance foundations) and must meet certain financial criteria. Uninsured patients and insured patients without coverage for their medicine must meet different financial criteria.

Please see full [Prescribing Information](#) and additional Important Safety Information throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1050 mg
INJECTION FOR IV USE

TECENTRIQ CO-PAY PROGRAM

TECENTRIQ Access Solutions can refer you to the TECENTRIQ Co-pay Program. It can help you with the out-of-pocket costs of TECENTRIQ, if you're eligible.

12

NO INCOME REQUIREMENTS	PATIENTS PAY AS LITTLE AS \$5 FOR TECENTRIQ	ANNUAL BENEFIT LIMIT OF CO-PAY PROGRAM IS \$25k
-------------------------------------	--	--

No physical card needed; patients simply need their Member ID.

To learn more about the TECENTRIQ Co-pay Program, call (855) MY-COPAY (855-692-6729) or visit copayassistancenow.com.

This TECENTRIQ Co-pay Program is valid ONLY for patients with commercial insurance who have a valid prescription for a Food and Drug Administration (FDA)-approved indication of a Genentech medication. Patients using Medicare, Medicaid, or any other federal or state government program to pay for their medications are not eligible.

Under the Program, the patient will pay a co-pay. After reaching the maximum Program benefit, the patient will be responsible for all out-of-pocket costs.

All participants are responsible for reporting the receipt of all Program benefits as required by any insurer or by law. No party may seek reimbursement for all or any part of the benefit received through this Program. This Program is void where prohibited by law. Genentech reserves the right to rescind, revoke, or amend the Program without notice at any time. Additional eligibility criteria apply. See full terms and conditions at copayassistancenow.com.

Please visit copayassistancenow.com for the full list of terms and conditions.


CONNECTING WITH CARE PARTNERS

13

There will be times that you need extra help and support. It's important to know that you don't have to go through this alone. Friends, family, social workers, and support groups are just some of the people around you that may help.

It may be hard to share your feelings with your caregiver, or to ask for help. Communication with your care partner is an important step in getting the support you need. The more they know how you're feeling, the better prepared they can be to help you.

Share more information and resources with your loved one by visiting TECENTRIQ.com/SCLC/Caregiver.



Please see full [Prescribing Information](#) and additional Important Safety Information throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE



TECENTRIO is not approved for children.

Caregiver Guide

Share the following pages with your loved ones. Helpful tips and advice are included as a guide for anyone in your support circle. Remember, you're not alone.

Please see full [Prescribing Information](#) and additional Important Safety Information throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE

CAREGIVERS

You play an important role in your loved one's treatment journey. Being a caregiver means that you may have to be there for a loved one who has been there for you. Your help can make this difficult time a little easier for your loved one. This brochure will provide you with information and resources that you may find useful.

Write down any questions in the notes section at the end of this brochure that you or your loved one would like to ask the healthcare team.

Below are some tips on how you may be able to help support your loved one throughout their journey:



Communication

- Talk to your loved one. Make sure they know you want to help and that communication is key (see the tips on how to communicate on page 18)
- Offer to go with your loved one to medical appointments
- Communicate with the entire healthcare team



Information Gathering

- As a caregiver, you can act as an extra set of eyes and ears. Ask important questions and gather information to share with your loved one
- Offer to help your loved one note any changes in symptoms or side effects
- Keep track of medical information and paperwork
- Find out if there are any special instructions and treatment tips that the healthcare team recommends



Emotional Support

- Try to keep a positive attitude-it is beneficial for your loved ones and for you
- Spend time together and do things that remind you both that you have a relationship separate from your loved one's diagnosis
- Encourage your loved one to engage in the activities they like as much as possible

TAKING CARE OF YOURSELF

You play an essential role in the TECENTRIQ journey. Taking care of yourself is an important step towards ensuring that you're taking care of your loved one.

17

Below are some tips that may help you avoid feeling overwhelmed:



- Keep a journal to organize thoughts and create memories
- Have regular doctor checkups
- Know your limits and be realistic about the time and energy you can give



- Find time to relax and do what you enjoy
- Go easy on yourself, focus on positive things you are doing for your loved one
- Consider routines that can improve your mood and energy



- Don't be afraid to ask for help; your family or friends can be your first resource
- Consider joining a support group or talking to a counselor



Please see full **Prescribing Information** and additional **Important Safety Information** throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1000 mg
INJECTION FOR IV USE

THE VALUE OF COMMUNICATION



Communication with your loved one is one of the most valuable tools you have.

The more you know about your loved one and how they're feeling, the better prepared you'll be to help them. Your support is valuable, and sometimes just being there to talk and listen can make a big difference.

Here are some different ways to help create a healthy dialogue:

- Ask if it's a good time to talk
- Be supportive, listen, and don't interrupt
- Think about what you want to say in advance
- Speak from the heart
- Remember that you can have more than one talk
- Don't always say, "It'll be okay"

Understanding your loved one's treatment is a helpful step towards keeping them on track.



IMPORTANT APPOINTMENTS

Please use this space to write down the dates and times of upcoming medical appointments that your loved one has, including TECENTRIQ infusion appointments.

Date: _____ **Time:** _____

Important notes: _____

Date: _____ **Time:** _____

Important notes: _____

Date: _____ **Time:** _____

Important notes: _____

Date: _____ **Time:** _____

Important notes: _____

IMPORTANT SAFETY INFORMATION ABOUT TECENTRIQ

What is the most important information about TECENTRIQ?

TECENTRIQ can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life threatening and can lead to death.

Call or see your healthcare provider right away if you get any symptoms of the following problems or these symptoms get worse.

TECENTRIQ can cause serious side effects, including:

- **Lung problems (pneumonitis)**—signs and symptoms of pneumonitis may include new or worsening cough, shortness of breath, and chest pain
- **Liver problems (hepatitis)**—signs and symptoms of hepatitis may include yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), drowsiness, dark urine (tea colored), bleeding or bruising more easily than normal, and feeling less hungry than usual
- **Intestinal problems (colitis)**—signs and symptoms of colitis may include diarrhea (loose stools) or more bowel movements than usual, blood or mucus in your stools or dark, tarry, sticky stools, and severe stomach area (abdomen) pain or tenderness
- **Hormone gland problems (especially the thyroid, adrenal glands, pancreas, and pituitary)**—signs and symptoms that your hormone glands are not working properly may include headaches that will not go away or unusual headaches, extreme tiredness, weight gain or weight loss, dizziness or fainting, feeling more hungry or thirsty than usual, hair loss, changes in mood or behavior (such as decreased sex drive, irritability, or forgetfulness), feeling cold, constipation, your voice gets deeper, urinating more often than usual, nausea or vomiting, and stomach area (abdomen) pain
- **Problems in other organs**—signs and symptoms may include severe muscle weakness, numbness or tingling in hands or feet, confusion, blurry vision, double vision, or other vision problems, changes in mood or behavior, extreme sensitivity to light, neck stiffness, eye pain or redness, skin blisters or peeling, chest pain, irregular heartbeat, shortness of breath, or swelling of the ankles
- **Severe infections**—signs and symptoms of infection may include fever, cough, flu-like symptoms, pain when urinating, and frequent urination or back pain
- **Severe infusion reactions**—signs and symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, swelling of your face or lips, dizziness, fever, feeling like passing out, and back or neck pain

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may delay or completely stop treatment with TECENTRIQ if you have severe side effects.

Before you receive TECENTRIQ, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems (such as Crohn's disease, ulcerative colitis, or lupus); have had an organ transplant; have lung or breathing problems; have liver problems; have a condition that affects your nervous system (such as myasthenia gravis or Guillain-Barre syndrome); or are being treated for an infection
- are pregnant or plan to become pregnant. TECENTRIQ can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TECENTRIQ. **Females who are able to become pregnant:**
 - your healthcare provider should do a pregnancy test before you start treatment with TECENTRIQ
 - you should use an effective method of birth control during your treatment and for at least 5 months after the last dose of TECENTRIQ
- are breastfeeding or plan to breastfeed. It is not known if TECENTRIQ passes into your breast milk. Do not breastfeed during treatment and for at least 5 months after the last dose of TECENTRIQ

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of TECENTRIQ when used in lung cancer with other anti-cancer medicines include:

- feeling tired or weak
- nausea
- hair loss
- constipation
- diarrhea
- decreased appetite

TECENTRIQ may cause fertility problems in females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TECENTRIQ. Ask your healthcare provider or pharmacist for more information about the benefits and side effects of TECENTRIQ.

You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at 1-888-835-2555.

Please see full **Prescribing Information** and additional **Important Safety Information** throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1000 mg
INJECTION FOR IV USE

GLOSSARY



Antibody:

A protein made by the immune system during an immune response that attaches to specific cells and can affect the immune system's ability to target and kill cells.

ES-SCLC (extensive-stage small cell lung cancer):

Cancer has spread outside of the lung in which it began or to other parts of the body.

Immune system:

A collection of organs, special cells, and substances that help protect you from infections and other diseases.

Infusion:

A method of putting fluids, including drugs, into the bloodstream. Also called intravenous infusion.

Median:

A statistics term. The middle value, or number, in a set of measurements, which have been put in order from lowest to highest.

PD-L1 (programmed death-ligand 1):

A protein found on the surface of many cells, including cancer cells, that can affect the immune system's ability to target and attack cells.

Protein:

A molecule that is the building block of the cells in the body. Proteins are needed for the body to function properly.

SCLC (small cell lung cancer):

An aggressive (fast-growing) cancer that forms in tissues of the lung and can spread to other parts of the body.

Sources:

<http://www.cancer.gov/publications/dictionaries/cancer-terms>.

<https://www.cancer.org/cancer/small-cell-lung-cancer/detection-diagnosis-staging/staging.html>.

The people depicted herein are not actual patients.

YOUR NOTES

23

Use the following pages to write down any notes, questions, or concerns you would like to mention to your healthcare team.



Please see full **Prescribing Information** and additional **Important Safety Information** throughout this brochure.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1000 mg
INJECTION FOR IV USE

Genentech is committed to advancing cancer treatment

Genentech has been committed to finding new ways of treating lung cancer.

A significant advancement in small cell lung cancer (SCLC) treatment, TECENTRIQ is approved as the first immunotherapy combination* for extensive-stage SCLC.

*TECENTRIQ + carboplatin + etoposide.

SELECT IMPORTANT SAFETY INFORMATION

TECENTRIQ can cause serious side effects, including:

- **Intestinal problems (colitis)**—signs and symptoms of colitis may include diarrhea (loose stools) or more bowel movements than usual, blood or mucus in your stools or dark, tarry, sticky stools, and severe stomach area (abdomen) pain or tenderness
- **Hormone gland problems (especially the thyroid, adrenal glands, pancreas, and pituitary)**—signs and symptoms that your hormone glands are not working properly may include headaches that will not go away or unusual headaches, extreme tiredness, weight gain or weight loss, dizziness or fainting, feeling more hungry or thirsty than usual, hair loss, changes in mood or behavior (such as decreased sex drive, irritability, or forgetfulness), feeling cold, constipation, your voice gets deeper, urinating more often than usual, nausea or vomiting, and stomach area (abdomen) pain

Please see pages 20-21 and full [Prescribing Information](#) for additional Important Safety Information.

Genentech
A Member of the Roche Group

© 2019 Genentech USA, Inc. All rights reserved.
PDL/111318/0213(1)

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE