

**phexxi™**  
(lactic acid, citric acid, and  
potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

**pH**INALLY! THERE'S **PHEXXI™**  
AN FDA-APPROVED, **HORMONE-FREE**  
BIRTH CONTROL GEL THAT YOU CAN  
USE IN THE MOMENT!

## 8 Simple Steps

Phexxi™ is a single-use product that is effective immediately, lasts up to 1 hour, and is only effective when used before sex. An additional dose of Phexxi™ must be applied for every act of vaginal sex (similar to using a new condom).

1



Wash your hands.

2




Remove the pre-filled applicator and plunger rod from the foil pouch.

3



Insert the plunger rod into the applicator, and push until the plunger rod connects to the applicator. Use a new pre-filled applicator if gel goes into the pink cap.

4



Remove the pink cap.

**PHEXXI™ IS NOW READY FOR USE!**

5



In an appropriate position, insert the Phexxi™ pre-filled applicator into your vagina as far as it will comfortably go while you continue to firmly hold it by the grooved area of the applicator.

6



Using your index finger, push the plunger rod until it stops to ensure you receive the entire dose of Phexxi™.

7



Remove the used Phexxi™ pre-filled applicator from your vagina. A small amount of gel may be left in the applicator, however, you will still get the right dose.

8



Throw away (dispose of) the used Phexxi™ pre-filled applicator and cap.

Avoid using Phexxi™ if you or your partner is allergic to any of the ingredients in Phexxi™.

## WHAT IS PHEXXI™?

Phexxi™ is a prescription vaginal gel used to prevent pregnancy in females who choose to use an on-demand method of birth control. Phexxi™ is only effective when used immediately **before** (or up to one hour before) each act of vaginal sex. Phexxi™ is not effective when used after vaginal sex.

**phexxi™**  
(lactic acid, citric acid, and  
potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

## IMPORTANT SAFETY INFORMATION

### WHAT ARE THE POSSIBLE SIDE EFFECTS OF PHEXXI™?

There have been a few cases of urinary bladder infection and kidney infection reported in clinical studies. One of the cases was serious. Avoid Phexxi™ if you have had repeated urinary tract infections or other urinary tract problems.

The **most common side effects** were vaginal burning, vaginal itching, vaginal yeast infection, urinary tract infection, vaginal area discomfort, bacterial vaginosis, and vaginal discharge. Women also reported genital discomfort, pain while urinating, and vaginal pain. Some male partners reported genital discomfort.

### WHAT ELSE SHOULD I KNOW ABOUT USING PHEXXI™?

Phexxi™ does not protect against any sexually transmitted diseases, including HIV. Avoid using Phexxi™ with a vaginal ring.

Contact your healthcare provider if you are experiencing severe genital irritation or discomfort or urinary tract symptoms. Avoid Phexxi™ if you or your sexual partner is allergic to lactic acid, citric acid, potassium bitartrate, or any of the ingredients in Phexxi™. Stop using Phexxi™ if you develop an allergic reaction.

**Please see full [Prescribing Information](#) for Phexxi™, including [Patient Information](#).**

***Please report side effects by contacting Evofem Biosciences® toll-free at 1-833-EVFMBIO or contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).***

Watch the full [Instructions for Use video](#)  
and learn more at [Phexxi.com](http://Phexxi.com)